


PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/127, 38/19		(11) International Publication Number: WO 99/27908
A1		(43) International Publication Date: 10 June 1999 (10.06.99)
(21) International Application Number: PCT/IL98/00586		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 1 December 1998 (01.12.98)		
(30) Priority Data: 60/067,697 4 December 1997 (04.12.97) US		
(71) Applicant (for all designated States except US): YISSUM RESEARCH DEVELOPMENT COMPANY OF THE HEBREW UNIVERSITY OF JERUSALEM [IL/IL]; Jabotinsky Street 46, 91042 Jerusalem (IL).		
(72) Inventors; and (75) Inventors/Applicants (for US only): GABIZON, Alberto, A. [IL/IL]; Bernstein Street 56/7, 96920 Jerusalem (IL). KEDAR, Eliezer [IL/IL]; Kadish Luz Street 24/7, 96920 Jerusalem (IL). BARENHOLZ, Yechezkel [IL/IL]; Nave Shanan Street 18, 93707 Jerusalem (IL).		
(74) Agent: REINHOLD COHN AND PARTNERS; P.O. Box 4060, 61040 Tel Aviv (IL).		

Published*With international search report.**Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.*

(54) Title: COMBINED CHEMO-IMMUNOTHERAPY WITH LIPOSOMAL DRUGS AND CYTOKINES



(57) Abstract

A method of antitumor therapy is described in which administration of a chemotherapeutic drug, encapsulated in liposomes, is supplemented by administration of an immunostimulating cytokine. The cytokine is preferably also encapsulated in liposomes. In tumor models for lung and colon carcinomas, this method produced a significantly greater therapeutic effect, as evidenced by survival rate and tumor size, than a combination of the effects produced by the free or liposome-encapsulated components administered individually.